



Amended Application for New License
Major Project – Existing Dam

Applicant-Prepared Draft
Essential Fish Habitat Assessment

Security Level: Public

Yuba River Development Project
FERC Project No. 2246

June 2017

©2017, Yuba County Water Agency
All Rights Reserved

Table of Contents
Description

Section No.	Description	Page No.
1.0	Introduction.....	1-1
1.1	Purpose of the Essential Fish Habitat Assessment	1-5
1.2	Regulatory Framework	1-5
1.2.1	Magnuson-Stevens Fishery Conservation and Management Act (16 U.S.C. §1801-1891d).....	1-5
2.0	Essential Fish Habitat Background.....	2-1
2.1	1999 Initial EFH Designation	2-4
2.2	2008 EFH Designation Codification.....	2-5
2.3	NMFS 2011 5-Year Review of EFH for Pacific Coast Salmon	2-5
2.4	Pacific Coast Salmon FMP Amendment 18	2-5
2.5	Pacific Coast Salmon FMP Amendment 19	2-6
3.0	Consultation History	3-1
3.1	EFH Consultations for the Proposed Action.....	3-1
3.2	Key Consultation Considerations	3-5
3.2.1	NMFS 2005 Biological Opinion (Yuba River Development Project License Amendment)	3-5
3.2.2	NMFS 2014 Biological Opinion (USACE’s Operation and Maintenance of Daguerre Point Dam)	3-6
3.2.3	NMFS 2014 Letter of Concurrence (USACE’s Operation and Maintenance of Englebright Dam).....	3-9
3.2.4	Other Activities	3-10
4.0	Description of the Proposed Action.....	4-1
4.1	Existing Project Facilities	4-1
4.1.1	New Colgate Development.....	4-2
4.1.2	Existing Project Boundary	4-9
4.1.3	Existing Project Safety.....	4-9
4.1.4	Existing Project Operations	4-9
4.1.5	Existing Environmental Measures	4-20
4.2	YCWA’s Proposal	4-29
4.2.1	Proposed Project Facilities.....	4-29
4.2.2	Proposed Project Boundary.....	4-63
4.2.3	Proposed Project Operations.....	4-67
4.2.4	Proposed Environmental Measures.....	4-74
5.0	Description of the EFH Action Area	5-1
5.1	NMFS Designation of EFH	5-1
5.2	FERC’s Defined Action Area	5-5

Table of Contents (continued)

Section No.	Description	Page No.
5.2.1	Yuba River Watershed Upstream of Englebright Dam and Reservoir	5-5
5.2.2	Yuba River Downstream of Englebright Dam and Reservoir	5-7
5.3	Downstream Extent of the Action Area.....	5-11
5.3.1	Feather River.....	5-16
5.3.2	Sacramento River.....	5-17
5.3.3	Sacramento-San Joaquin Delta	5-17
5.3.4	Pacific Ocean	5-18
6.0	Species Managed under the Pacific Coast Salmon Fisheries Management Plan.....	6-1
6.1	Central Valley Fall-/Late Fall-run Chinook Salmon ESU	6-1
6.1.1	General Life History and Habitat Requirements	6-2
6.1.2	Historical Distribution and Abundance	6-6
6.1.3	Population Status and Trends.....	6-10
6.2	Limiting Factors, Threats and Stressors.....	6-15
6.2.1	Pacific Coast Salmon	6-15
6.2.2	Central Valley Chinook Salmon	6-17
6.2.3	Yuba River Chinook Salmon	6-20
7.0	Overview of Chinook Salmon EFH Components in the Action Area	7-1
7.1	Yuba River Watershed Upstream of Englebright Dam	7-1
7.2	Yuba River Downstream of Englebright Dam.....	7-2
7.3	Migratory Habitat (Adult Upstream and Juvenile Downstream).....	7-3
7.3.1	Yuba River Watershed Upstream of Englebright Dam	7-4
7.3.2	Downstream of Englebright Dam	7-22
7.4	Spawning and Embryo Incubation Habitat	7-42
7.4.1	Yuba River Watershed Upstream of Englebright Dam	7-43
7.4.2	Downstream of Englebright Dam	7-50
7.5	Juvenile Rearing Habitat.....	7-79
7.5.1	Complex Channels and Floodplain Habitats.....	7-79
7.5.2	Thermal Refugia	7-80
7.5.3	Yuba River Watershed Upstream of Englebright Dam	7-81
7.5.4	Downstream of Englebright Dam	7-90
8.0	Analyses of Potential Effects of the Proposed Action on Essential Fish Habitat for Managed Species	8-1
8.1	EFH Assessment Approach	8-1
8.1.1	Habitat Descriptors	8-2

Table of Contents (continued)

Section No.	Description	Page No.
8.2	Summary of Existing EFH Conditions (Environmental Baseline Condition)	8-3
8.2.1	Yuba River Watershed Upstream of Englebright Dam	8-3
8.2.2	Yuba River Downstream of Englebright Dam.....	8-20
8.3	Potential Effects of the Proposed Action on EFH in the Action Area.....	8-27
8.3.1	Yuba River Watershed Upstream of Englebright Dam	8-30
8.3.2	Yuba River Downstream of Englebright Dam.....	8-55
8.4	Past, Present, and Reasonably Foreseeable Future Actions that Potentially Affect EFH in the Action Area (Cumulative Effects)	8-90
8.4.1	Past and Present Actions	8-91
8.4.2	Reasonably Foreseeable Future Actions	8-92
8.4.3	Yuba River Watershed Upstream of Englebright Dam	8-95
8.4.4	Yuba River Downstream of Englebright Dam.....	8-105
8.4.5	Other Future Activities in the Lower Yuba River	8-136
8.4.6	Out-of-Basin Future Activities	8-148
9.0	Summary of Effects	9-1
9.1	Summary of Anticipated Effects of the Proposed Action on EFH in the Action Area.....	9-1
9.1.1	Yuba River Watershed Upstream of Englebright Dam	9-1
9.1.2	Yuba River Downstream of Englebright Dam.....	9-3
9.2	Summary of Anticipated Effects of the Cumulative Condition on EFH in the Action Area.....	9-4
9.2.1	Yuba River Watershed Upstream of Englebright Dam	9-4
9.2.2	Yuba River Downstream of Englebright Dam.....	9-5
10.0	Determination	10-1
11.0	References.....	11-1

List of Figures

Figure No.	Description	Page No.
3.2-1.	Yuba Accord North Yuba Index.	3-11
4.1-1.	Our House Diversion Dam features.....	4-2
4.1-2.	Log Cabin Diversion Dam features.	4-3
4.1-3.	Longitudinal schematic of New Bullards Bar Dam on the North Yuba River and New Colgate Power Tunnel and Powerhouse on the Yuba River.....	4-6
4.1-4.	Longitudinal schematic of Narrows 2 Powerhouse Penstock and Powerhouse.....	4-8

List of Figures (continued)
Description

Figure No.	Description	Page No.
4.1-5.	Historical mean daily discharge from Narrows 1 Powerhouse, Narrows 2 facilities, the Smartsville gage, and Englebright Dam spill in Water Year 2007. The Narrows 2 Powerhouse was shut down during the period of October 1 through December 30, 2006 for the installation of the Narrows 2 Full Bypass. Siphons over Englebright Dam were used to provide flow to the Yuba River in addition to Narrows 1 Powerhouse releases. Accordingly, there is a difference between the flow from the Narrows 1 Powerhouse and Smartsville gage.	4-15
4.1-6.	Historical mean daily discharge from Narrows 1 Powerhouse, Narrows 2 facilities, the Smartsville gage, and Englebright Dam spill in Water Year 2008.....	4-16
4.1-7.	Historical mean daily discharge from Narrows 1 Powerhouse, Narrows 2 facilities, the Smartsville gage, and Englebright Dam spill in Water Year 2009.....	4-16
4.1-8.	Historical mean daily discharge from Narrows 1 Powerhouse, Narrows 2 facilities, the Smartsville gage, and Englebright Dam spill in Water Year 2010.....	4-17
4.1-9.	Historical mean daily discharge from Narrows 1 Powerhouse, Narrows 2 facilities, the Smartsville gage, and Englebright Dam spill in Water Year 2011.....	4-17
4.1-10.	Historical mean daily discharge from Narrows 1 Powerhouse, Narrows 2 facilities, the Smartsville gage, and Englebright Dam spill in Water Year 2012.....	4-18
4.1-11.	Historical mean daily discharge from Narrows 1 Powerhouse, Narrows 2 facilities, the Smartsville gage, and Englebright Dam spill in Water Year 2013.....	4-18
4.1-12.	Historical mean daily discharge from Narrows 1 Powerhouse, Narrows 2 facilities, the Smartsville gage, and Englebright Dam spill in Water Year 2014.....	4-19
4.1-13.	Historical mean daily discharge from Narrows 1 Powerhouse, Narrows 2 facilities, the Smartsville gage, and Englebright Dam spill in Water Year 2015.....	4-19
4.1-14.	Historical mean daily discharge from Narrows 1 Powerhouse, Narrows 2 facilities, the Smartsville gage, and Englebright Dam spill in Water Year 2016.....	4-20
4.1-15.	YCWA’s Member Unit service areas.	4-26
4.2-1.	Concept-level plan schematic flow diagram for YCWA’s proposed New Colgate Powerhouse TDS.....	4-31
4.2-2.	Concept-level plan for general arrangement for YCWA’s proposed New Colgate Powerhouse TDS.....	4-32

List of Figures (continued)
Description

Figure No.	Description	Page No.
4.2-3.	Concept-level plan for piping sections and details for YCWA’s proposed New Colgate Powerhouse TDS.	4-33
4.2-4.	Concept-level plan and profile for YCWA’s proposed New Bullards Bar Dam Auxiliary Flood Control Outlet.	4-39
4.2-5.	Concept-level map of construction area for YCWA’s proposed New Bullards Bar Auxiliary Flood Control Outlet.	4-43
4.2-6.	Concept-level profile of YCWA’s proposed Our House Diversion Dam Fish Release Outlet.	4-45
4.2-7.	Concept-level profile of YCWA’s proposed Log Cabin Diversion Dam Fish Release Outlet.	4-46
4.2-8.	Concept-level details of YCWA’s proposed Lohman Ridge Inlet Control Gate and Debris Rake.	4-49
5.1-1.	Chinook salmon EFH in California (PFMC 2014).	5-4
5.2-2.	Map of study area showing geomorphic reach boundaries.	5-9
5.2-3.	Map of habitat/hydrologic zone boundaries.	5-10
5.3-1.	Other EFH areas affecting Yuba River spring- and fall-run Chinook salmon throughout the ESUs.	5-15
7.3-1.	Mean daily water temperatures in the North Yuba River downstream of New Bullards Bar Dam (RM 22.4) and upstream of the Middle Yuba River confluence (RM 0.1) in WYs 2009, 2010, 2011 and 2012.	7-9
7.3-2.	Mean daily water temperatures in the Middle Yuba River from Our House Diversion Dam to the North Yuba River confluence during WYs 2009, 2010, 2011 and 2012.	7-16
7.3-3.	Mean daily water temperatures in the Yuba River from the Middle and North Yuba rivers downstream to above the normal-maximum water-surface elevation of the USACE’s Englebright Reservoir during WYs 2009, 2010, 2011 and 2012.	7-20
7.3-4.	Locations of Narrows 2 Powerhouse, Narrows 2 Partial Bypass, Narrows 2 Full Bypass and Narrows 1 Powerhouse on the Yuba River in relation to Englebright Dam. In this photo, flow is being released from the Narrows 1 and 2 powerhouses, and no flow is being released from the Narrows 2 Partial Bypass or the Full Bypass.	7-24
7.3-5.	Monitored lower Yuba River water temperatures and spring-run Chinook salmon upper tolerable water temperature index values.	7-34
7.3-6.	Monitored lower Yuba River water temperatures and fall-run Chinook salmon upper tolerable water temperature index values.	7-35
7.4-1.	North Yuba River – New Bullards Bar Dam Reach – Monthly habitat exceedance results shown as AUC for rainbow trout spawning – Node 0.	7-44

List of Figures (continued)
Description

Figure No.	Description	Page No.
7.4-2.	Middle Yuba River – Oregon Creek Reach – Monthly habitat exceedance results shown as AUC for rainbow trout spawning.	7-47
7.4-3.	Yuba River – Middle/North Yuba Reach – Monthly habitat exceedance results shown as AUC for rainbow trout spawning.	7-49
7.4-4.	Yuba River – New Colgate Powerhouse Reach – Monthly habitat exceedance results shown as AUC for rainbow trout spawning.	7-49
7.4-5.	Longitudinal distribution of the mean substrate diameter. The box represents the typical range of spawning substrate sizes observed on the lower Yuba River.	7-53
7.4-6.	Spring-run Chinook salmon spawning habitat duration over the 41-year hydrologic period for the Environmental Baseline and Without-Project scenarios.	7-56
7.4-7.	Fall-run Chinook salmon spawning habitat duration over the 41-year hydrologic period for the Environmental Baseline and Without-Project scenarios.	7-58
7.4-8.	Diagrammatic side view of a Chinook salmon redd showing the relative location and mean depth of egg pockets on the Trinity River, CA, as reported in Evenson (2001).	7-60
7.4-9.	Cumulative proportions of spring-run Chinook salmon newly-built redds observed in lower Yuba River reaches upstream of Daguerre Point Dam during the 2009 and 2010 redd surveys.1	7-62
7.4-10.	Daily distribution of spawning temporal weighting coefficients (wd) used in the spring-run Chinook salmon redd dewatering indices.	7-63
7.4-11.	Cumulative proportions of Chinook salmon newly-built redds observed in lower Yuba River reaches upstream and downstream Daguerre Point Dam during the 2009 and 2010 redd surveys (circles and squares) and corresponding fitted asymmetric logistic curves used to derive the temporal weighting coefficients for the fall-run Chinook salmon redd dewatering indices.	7-66
7.4-12.	Distributions of daily temporal weighting coefficients used in the calculation of redd dewatering indices for fall-run Chinook salmon spawning upstream (green bars, wd U) and downstream (orange bars, wd D) of Daguerre Point Dam. The distribution of daily temporal weighting coefficients for the combined reach is described by the gray line (wd = wd U + wd D).	7-67
7.4-13.	Cumulative proportions of redd depths measured in the lower Yuba River during the 2011 weekly Chinook salmon redd surveys, and derived asymmetric logistic curve used to assess the expected redd depths of both spring-run and fall-run Chinook salmon spawning in the lower Yuba River.	7-69

List of Figures (continued)
Description

Figure No.	Description	Page No.
7.4-14.	Cumulative proportions of egg depths derived from the 2011 weekly Chinook salmon redd surveys in the lower Yuba River, and derived asymmetric logistic curve used to assess the expected depths of the egg pockets for both spring-run and fall-run Chinook salmon spawning in the lower Yuba River.....	7-70
7.5-1.	Overview of scores by basin, stream, and indices. Sites with starred symbols represent locations where insufficient organisms were collected to make the resultant IBI and MMI scores reliable.....	7-86
7.5-2.	Spring-run Chinook salmon fry in-channel rearing habitat duration over the 41-year hydrologic period for the Environmental Baseline and Without-Project scenarios.....	7-94
7.5-3.	Spring-run Chinook salmon juvenile in-channel rearing habitat duration over the 41-year hydrologic period for the Environmental Baseline and Without-Project scenarios.....	7-96
7.5-4.	Fall-run Chinook salmon fry in-channel rearing habitat duration over the 41-year hydrologic period for the Environmental Baseline and Without-Project scenarios.	7-98
7.5-5.	Fall-run Chinook salmon juvenile in-channel rearing habitat duration over the 41-year hydrologic period for the Environmental Baseline and Without-Project scenarios.....	7-100
7.5-6.	Comparison of the amount (acres) of spring-run Chinook salmon fry weighted usable area (WUA) without cover under the Environmental Baseline and the Without-Project scenarios over the 41-year period of evaluation. Shown are the amounts over the long-term full simulation period (all years) and by water year type of total habitat provided on days when flows were $\leq 5,000$ cfs and for days when flows were $> 5,000$ cfs.	7-102
7.5-7.	Comparison of the amount (acres) of spring-run Chinook salmon juvenile weighted usable area (WUA) without cover under the Environmental Baseline and the Without-Project scenarios over the 41-year period of evaluation. Shown are the amounts over the long-term full simulation period (all years) and by water year type of total habitat provided on days when flows were $\leq 5,000$ cfs and for days when flows were $> 5,000$ cfs.	7-104
7.5-8.	Comparison of the amount (acres) of fall-run Chinook salmon fry weighted usable area (WUA) without cover under the Environmental Baseline and the Without-Project scenarios over the 41-year period of evaluation. Shown are the amounts over the long-term full simulation period (all years) and by water year type of total habitat provided on days when flows were $\leq 5,000$ cfs and for days when flows were $> 5,000$ cfs.	7-106

List of Figures (continued)
Description

Figure No.	Description	Page No.
7.5-9.	Comparison of the amount (acres) of fall-run Chinook salmon juvenile weighted usable area (WUA) without cover under the Environmental Baseline and the Without-Project scenarios over the 41-year period of evaluation. Shown are the amounts over the long-term full simulation period (all years) and by water year type of total habitat provided on days when flows were $\leq 5,000$ cfs and for days when flows were $> 5,000$ cfs.	7-108
7.5-10.	Vegetation communities in the lower Yuba River vicinity.	7-111
7.5-11.	Average percent of all off-channel areas in the lower Yuba River experiencing the specified number of isolation events over the 41-year hydrologic period for the Environmental Baseline and Without-Project scenarios.....	7-119
8.3-1.	Spring-run Chinook salmon spawning habitat duration over the 41-year hydrologic period for the Proposed Action and Environmental Baseline.	8-58
8.3-2.	Fall-run Chinook salmon spawning habitat duration over the 41-year hydrologic period for the Proposed Action and Environmental Baseline.	8-60
8.3-3.	Spring-run Chinook salmon fry in-channel rearing habitat duration over the 41-year hydrologic period for the Proposed Action and Environmental Baseline.....	8-66
8.3-4.	Fall-run Chinook salmon fry in-channel rearing habitat duration over the 41-year hydrologic period for the Proposed Action and Environmental Baseline.....	8-68
8.3-5.	Spring-run Chinook salmon juvenile in-channel rearing habitat duration over the 41-year hydrologic period for the Proposed Action and Environmental Baseline.....	8-70
8.3-6.	Fall-run Chinook salmon juvenile in-channel rearing habitat duration over the 41-year hydrologic period for the Proposed Action and Environmental Baseline.....	8-72
8.3-7.	Comparison of the amounts (in acres) of spring-run Chinook salmon fry weighted usable area (WUA) without cover under the Proposed Action and the Environmental Baseline over the 41-year period of evaluation. Shown are the amounts over the long-term full simulation period (all years) and by water year type of total habitat provided on days when flows were $\leq 5,000$ cfs and for days when flows were $> 5,000$ cfs.....	8-74
8.3-8.	Comparison of the amounts (in acres) of fall-run Chinook salmon fry weighted usable area (WUA) without cover under the Proposed Action and the Environmental Baseline over the 41-year period of evaluation. Shown are the amounts over the long-term full simulation period (all years) and by water year type of total habitat provided on days when flows were $\leq 5,000$ cfs and for days when flows were $> 5,000$ cfs.....	8-76

List of Figures (continued)
Description

Figure No.	Description	Page No.
8.3-9.	Comparison of the amounts (in acres) of spring-run Chinook salmon juvenile weighted usable area (WUA) without cover under the Proposed Action and the Environmental Baseline over the 41-year period of evaluation. Shown are the amounts over the long-term full simulation period (all years) and by water year type of total habitat provided on days when flows were $\leq 5,000$ cfs and for days when flows were $> 5,000$ cfs.	8-78
8.3-10.	Comparison of the amounts (in acres) of fall-run Chinook salmon juvenile weighted usable area (WUA) without cover under the Proposed Action and the Environmental Baseline over the 41-year period of evaluation. Shown are the amounts over the long-term full simulation period (all years) and by water year type of total habitat provided on days when flows were $\leq 5,000$ cfs and for days when flows were $> 5,000$ cfs.	8-80
8.3-11.	Average percent of all off-channel areas in the lower Yuba River experiencing the specified number of isolation events over the 41-year hydrologic period for the Proposed Action and Environmental Baseline.	8-82
8.3-12.	Exceedance probability of specified daily stage reductions (cm/day) from April 1 through July 15 under the Proposed Action and the Environmental baseline over the 41-year period of evaluation.	8-89
8.4-1.	Spring-run Chinook salmon spawning habitat duration over the 33-year hydrologic period for the Cumulative Condition and Environmental Baseline.	8-108
8.4-2.	Fall-run Chinook salmon spawning habitat duration over the 33-year hydrologic period for the Cumulative Condition and Environmental Baseline.	8-110
8.4-3.	Spring-run Chinook salmon fry in-channel rearing habitat duration over the 33-year hydrologic period for the Cumulative Condition and Environmental Baseline.	8-115
8.4-4.	Comparison of the amount (in acres) of spring-run Chinook salmon fry weighted usable area (WUA) without cover under the Cumulative Condition and the Environmental Baseline over the 33-year period of evaluation. Shown are the amounts over the long-term full simulation period (all years) and by water year type of total habitat provided on days when flows were $\leq 5,000$ cfs and for days when flows were $> 5,000$ cfs.	8-117
8.4-5.	Spring-run Chinook salmon juvenile in-channel rearing habitat duration over the 33-year hydrologic period for the Cumulative Condition and Environmental Baseline.	8-118

List of Figures (continued)

Figure No.	Description	Page No.
8.4-6.	Comparison of the amounts (in acres) of spring-run Chinook salmon juvenile weighted usable area (WUA) without cover under the Cumulative Condition and the Environmental Baseline over the 33-year period of evaluation. Shown are the amounts over the long-term full simulation period (all years) and by water year type of total habitat provided on days when flows were $\leq 5,000$ cfs and for days when flows were $> 5,000$ cfs.	8-120
8.4-7.	Fall-run Chinook salmon fry in-channel rearing habitat duration over the 33-year hydrologic period for the Cumulative Condition and Environmental Baseline.	8-121
8.4-8.	Comparison of the amounts (in acres) of fall-run Chinook salmon fry weighted usable area (WUA) without cover under the Cumulative Condition and the Environmental Baseline over the 33-year period of evaluation. Shown are the amounts over the long-term full simulation period (all years) and by water year type of total habitat provided on days when flows were $\leq 5,000$ cfs and for days when flows were $> 5,000$ cfs.	8-123
8.4-9.	Fall-run Chinook salmon juvenile in-channel rearing habitat duration over the 33-year hydrologic period for the Cumulative Condition and Environmental Baseline.	8-125
8.4-10.	Comparison of the amounts (in acres) of fall-run Chinook salmon juvenile weighted usable area (WUA) without cover under the Cumulative Condition and the Environmental Baseline over the 33-year period of evaluation. Shown are the amounts over the long-term full simulation period (all years) and by water year type of total habitat provided on days when flows were $\leq 5,000$ cfs and for days when flows were $> 5,000$ cfs.	8-127
8.4-11.	Average percent of all off-channel areas in the lower Yuba River experiencing the specified number of isolation events over the 33-year hydrologic period for the Cumulative Condition and Environmental Baseline.	8-129
8.4-12.	Exceedance probability of specified daily stage reductions (cm/day) from April 1 through July 15 under the Cumulative Condition and the Environmental baseline over the 33-year period of evaluation.	8-135
8.4-13.	Aggregate mines in the Yuba Goldfields (Source: SMGB 2014).	8-139
8.4-14.	Conceptual simulation reclaimed areas in the Yuba Goldfields resulting from implementation of Western Aggregate’s Amended Reclamation Plan (SMGB 2014).	8-142

List of Tables

Table No.	Description	Page No.
2.0-1.	List of 10(j) potential recommendations on which YCWA, USFWS and Cal Fish and Wildlife have reached agreement.	2-3

List of Tables (continued)

Table No.	Description	Page No.
3.2-1.	Yuba Accord flow schedules.	3-11
4.1-1.	Key information regarding Yuba River Development Project powerhouses.	4-1
4.1-2.	Key information regarding Yuba River Development Project reservoirs and impoundments.	4-1
4.1-3.	Typical distribution of flows under normal operations (i.e., excluding brief transition periods) among Narrows 2 Powerhouse (generation only), Partial Bypass, Full Bypass and Narrows 1 Powerhouse.	4-14
4.1-4.	New Bullards Bar Reservoir flood storage space allocation in thousands of ac-ft.	4-23
4.1-5.	Water right licenses held by YCWA for operation of the Project for power generation.	4-24
4.1-6.	YCWA’s annual contract amounts and places of delivery.	4-25
4.1-7.	YCWA’s water supply contract shortage provisions.	4-27
4.1-8.	YCWA historical sales from 1987 to 2016.	4-28
4.2-1.	Primary Project Roads and Trails (non-recreation roads and trails) included in the Yuba River Development Project.	4-60
4.2-2.	Recreation roads included in the Yuba River Development Project.	4-61
4.2-3.	Proposed additions to Primary Project Roads and Trails (non-recreation roads and trails) included in Yuba River Development Project by Development.	4-62
4.2-4.	Proposed additions to and withdrawals of Recreation Roads associated with the New Colgate Development. YCWA does not propose any changes to Recreation Roads and Trails associated with the New Bullards Bar Minimum Flow Development or the Narrows 2 Development.	4-62
4.2-5.	Summary of land ownership within the proposed Yuba River Development Project FERC Project Boundary by Project Development and difference as compared to existing FERC Project Boundary.	4-67
4.2-6.	Smartsville hydrological index Water Year Types and associated thresholds.	4-68
4.2-7.	North Yuba Index1 Schedules and associated thresholds.	4-68
4.2-8.	Proposed Project flow requirements for the Middle Yuba River downstream of Our House Diversion Dam by Smartsville hydrological index Water Year Type.	4-69
4.2-9.	Proposed Project flow requirements for Oregon Creek downstream of Log Cabin Diversion Dam by Smartsville Hydrological Index Water Year Type.	4-69

List of Tables (continued)
Description

Table No.	Description	Page No.
4.2-10.	Proposed Project Flow Requirements for the North Yuba River downstream of New Bullards Bar Dam by Smartsville Hydrological Index Water Year Type.....	4-70
4.2-11.	Proposed Project whitewater boating flows below Our House Diversion Dam.....	4-71
4.2-12.	Proposed Project flow requirements for the Yuba River downstream of Narrows 2 Powerhouse and Narrows 2 Full Flow Bypass by North Yuba Index Flow Schedule.....	4-71
4.2-13.	Maximum flow reductions corresponding to the maximum 5-day average release (Base Flow) that has occurred during the period extending from September 1 through December 31.....	4-73
4.2-14.	Maximum flow reductions corresponding to the maximum 5-day average release (Base Flow) that has occurred during the period extending from January 1 through May 31.	4-73
4.2-15.	Maximum flow reductions corresponding to the preceding day average flow that has occurred during the period extending from April 1 through July 15.....	4-73
4.2-16.	List of YCWA’s proposed conditions and the Relicensing Participants that YCWA understands agree with YCWA’s proposed conditions, as provided in Appendix E2 of the Amended FLA. A green shaded row indicates a condition that YCWA and the Forest Service have reached tentative agreement on as a FPA § 4(e) potential condition. A yellow shaded row indicates a condition that YCWA and Cal Fish and Wildlife or USFWS have reached tentative agreement on as a FPA § 10(j) potential recommendation. A white shaded row indicates a condition that has not been collaboratively agreed to between YCWA and agencies.	4-75
5.1-1.	Pacific Salmon EFH Identified by USGS Hydrologic Unit Code (HUC) for the Yuba River (73 FR 60987).....	5-1
5.2-1.	Geomorphic reaches in the Yuba River downstream from Englebright Dam.....	5-8
5.2-2.	Hydrologic zone boundaries established for habitat modeling in the Yuba River downstream from Englebright Dam.....	5-8
6.1-1.	Lifestage-specific periodicities for fall-run Chinook salmon in the lower Yuba River.....	6-2
6.1-2.	Fall-run Chinook salmon escapement estimates from 1994-2015 for rivers in the California Central Valley (CDFW 2016).....	6-8
6.1-3.	Fall-run Chinook salmon abundance from 1996 through 2015 in the Sacramento River system.....	6-11

List of Tables (continued)
Description

Table No.	Description	Page No.
6.1-4.	Estimated number of phenotypic spring-run and fall-run Chinook salmon spawning in the lower Yuba River from 2004 through 2015, and corresponding spring-run and fall-run Chinook salmon spawners reported in GrandTab for Sacramento River system.....	6-14
6.2-1.	Lifestage-specific habitat concerns associated with Pacific Coast salmon EFH (PFMC 1999).....	6-16
6.2-2.	Non-fishing threats to Pacific Coast salmon EFH. Newly identified threats appear in the right column. Detailed information on the threats identified in the first column can be found in Amendment 14.....	6-16
7.3-1.	Length, frequency, width and depth of ground-mapped habitat units for the North Yuba River – New Bullards Bar Reach (between the junction with the Middle Yuba River and New Bullards Bar Dam). The shaded cells are characteristics of pools that do not apply to non-pool habitat types.....	7-5
7.3-2.	Summary of ground mapped data for the North Yuba River – New Bullards Bar Reach (between the junction with the Middle Yuba River and New Bullards Bar Dam).....	7-5
7.3-3.	Summary of minimum and maximum daily average water temperatures (°C) by month in the upper Yuba River watershed potentially affected by the Proposed Action during WYs 2009 through 2012. Shaded cells indicate values over 20°C (68°F).....	7-7
7.3-4.	Difference in simulated upper tolerable water temperature exceedance probabilities for spring-run Chinook salmon lifestages under the Environmental Baseline scenario, relative to the Without-Project scenario.....	7-12
7.3-5.	Difference in simulated upper tolerable water temperature exceedance probabilities for fall-run Chinook salmon lifestages under the Environmental Baseline scenario, relative to the Without-Project scenario.....	7-12
7.3-6.	Length, frequency, width and depth of ground-mapped habitat units in the Middle Yuba River – Oregon Creek and Our House Diversion Dam Reaches (between the junction with North Yuba River to Our House Diversion Dam). The shaded cells are characteristics of pools that do not apply to non-pool habitat types.....	7-15
7.3-7.	Reach summary of ground mapped data for the Middle Yuba River – Oregon Creek and Our House Diversion Dam Reaches (between the junction with North Yuba River to Our House Diversion Dam).....	7-15
7.3-8.	Length, frequency, width and depth of ground-mapped habitat units for the Mainstem Yuba River – New Colgate Powerhouse and Middle/North Yuba River Reaches (between the New Colgate Powerhouse and the Middle/North Yuba junction). The shaded cells are characteristics of pools that do not apply to non-pool habitat types.....	7-19

List of Tables (continued)

Table No.	Description	Page No.
7.3-9.	Summary of ground mapped data for the Mainstem Yuba River – New Colgate Powerhouse and Middle/North Yuba River Reaches (between the New Colgate Powerhouse and the Middle/North Yuba junction).	7-19
7.3-10.	Spring-run Chinook salmon lifestage-specific upper tolerable water temperature index values.	7-32
7.3-11.	Fall-run Chinook salmon lifestage-specific upper tolerable water temperature index values.	7-33
7.3-12.	Difference in simulated upper tolerable water temperature exceedance probabilities for spring-run Chinook salmon lifestages under the Environmental Baseline, relative to the Without-Project scenario.	7-37
7.3-13.	Difference in simulated upper tolerable water temperature exceedance probabilities for fall-run Chinook salmon lifestages under the Environmental Baseline, relative to the Without-Project scenario.	7-37
7.4-1.	Difference in simulated upper tolerable water temperature exceedance probabilities for spring-run Chinook salmon lifestages under the Environmental Baseline scenario, relative to the Without-Project scenario.	7-45
7.4-2.	Difference in simulated upper tolerable water temperature exceedance probabilities for fall-run Chinook salmon lifestages under the Environmental Baseline scenario, relative to the Without-Project scenario.	7-45
7.4-3.	Long-term and WYT average spring-run Chinook salmon spawning WUA (percent of maximum) under the Environmental Baseline and Without-Project scenarios.	7-55
7.4-4.	Long-term and WYT average fall-run Chinook salmon spawning WUA (percent of maximum) under the Environmental Baseline and Without-Project scenarios.	7-57
7.4-5.	Number and proportion of newly-built Chinook salmon redds observed in the reaches upstream of Daguerre Point Dam during the 2009 and 2010 weekly redd surveys used to derive the temporal weighting coefficients (wd) for the spring-run Chinook salmon redd dewatering indices.	7-61
7.4-6.	Number and proportion of newly-built Chinook salmon redds observed in morphological units located upstream of Daguerre Point Dam during the 2009 and 2010 weekly redd surveys used as the spatial weighting coefficients (wh).	7-64
7.4-7.	Number and proportion of newly-built Chinook salmon redds observed in the reaches upstream of Daguerre Point Dam during the 2009 and 2010 weekly redd surveys used to derive the temporal weighting coefficients (wdU and wdD) for the fall-run Chinook salmon redd dewatering indices.	7-65

List of Tables (continued)
Description

Table No.	Description	Page No.
7.4-8.	Number and proportion of newly-built Chinook salmon redds observed in morphological units located upstream (UP) and downstream (DOWN) of Daguerre Point Dam during the 2009 and 2010 weekly redd surveys used in the calculation of the spatial weighting coefficients $w_h U$ and $w_h D$ in the fall-run Chinook salmon redd dewatering indices.	7-68
7.4-9.	Look-up stage-discharge table for Chinook salmon redds in morphological units located upstream of Daguerre Point Dam. Yellow highlighted cells contain stages calculated using the procedure described in text.	7-72
7.4-10.	Look-up stage-discharge table for Chinook salmon redds in morphological units located downstream of Daguerre Point Dam. Yellow highlighted cells contain stages calculated using the procedure described in text.	7-72
7.4-11.	Estimated spring-run Chinook salmon redd and egg pocket potential dewatering under the Environmental Baseline scenario relative to the Without-Project scenario.	7-77
7.4-12.	Estimated fall-run Chinook salmon redd and egg pocket potential dewatering under the Environmental Baseline scenario relative to the Without-Project scenario.	7-78
7.5-1.	Difference in simulated upper tolerable water temperature exceedance probabilities for spring-run Chinook salmon lifestages under the Environmental Baseline scenario, relative to the Without-Project scenario.....	7-84
7.5-2.	Difference in simulated upper tolerable water temperature exceedance probabilities for fall-run Chinook salmon lifestages under the Environmental Baseline scenario, relative to the Without-Project scenario.....	7-84
7.5-3.	Long-term and WYT average spring-run Chinook salmon fry in-channel rearing WUA (percent of maximum) under the Environmental Baseline and Without-Project scenarios.	7-93
7.5-4.	Long-term and WYT average spring-run Chinook salmon juvenile in-channel rearing WUA (percent of maximum) under the Environmental Baseline and Without-Project scenarios.	7-95
7.5-5.	Long-term and WYT average fall-run Chinook salmon fry in-channel rearing WUA (percent of maximum) under the Environmental Baseline and Without-Project scenarios.	7-97
7.5-6.	Long-term and WYT average fall-run Chinook salmon juvenile in-channel rearing WUA (percent of maximum) under the Environmental Baseline and Without-Project scenarios.	7-99
7.5-7.	Acres of spring-run Chinook salmon fry weighted usable area (WUA) without cover under the Environmental Baseline and the Without-Project scenarios over the 41-year period of evaluation for days when flows were $\leq 5,000$ cfs and for days when flows were $> 5,000$ cfs, and the differences between the two scenarios over the long-term full simulation period and by water year type.	7-101

List of Tables (continued)

Table No.	Description	Page No.
7.5-8.	Acres of spring-run Chinook salmon juvenile weighted usable area (WUA) without cover under the Environmental Baseline and the Without-Project scenarios over the 41-year period of evaluation for days when flows were $\leq 5,000$ cfs and for days when flows were $> 5,000$ cfs, and the differences between the two scenarios over the long-term full simulation period and by water year type.	7-103
7.5-9.	Acres of fall-run Chinook salmon fry weighted usable area (WUA) without cover under the Environmental Baseline and the Without-Project scenarios over the 41-year period of evaluation for days when flows were $\leq 5,000$ cfs and for days when flows were $> 5,000$ cfs, and the differences between the two scenarios over the long-term full simulation period and by water year type.	7-105
7.5-10.	Acres of fall-run Chinook salmon juvenile weighted usable area (WUA) without cover under the Environmental Baseline and the Without-Project scenarios over the 41-year period of evaluation for days when flows were $\leq 5,000$ cfs and for days when flows were $> 5,000$ cfs, and the differences between the two scenarios over the long-term full simulation period and by water year type.	7-107
8.2-1.	Simulated median monthly Environmental Baseline and Without-Project flows for October 1969 through September 2010 for the North Yuba River downstream of New Bullards Bar Dam by WYT.	8-5
8.2-2.	Simulated median monthly Environmental Baseline and Without-Project flows for October 1969 through September 2010 for the Middle Yuba River upstream of the North Yuba River confluence by WYT.	8-10
8.2-3.	Simulated median monthly Environmental Baseline and Without-Project flows for October 1969 through September 2010 for the Yuba River downstream of the Middle and North Yuba River confluence by WYT.	8-16
8.2-4.	Simulated median monthly Environmental Baseline and Without-Project flows for October 1969 through September 2010 for the Yuba River downstream of the New Colgate Powerhouse by WYT.	8-17
8.2-5.	Chinook salmon stressors and associated magnitudes in the lower Yuba River under the Environmental Baseline.	8-21
8.3-1.	Flows in the North Yuba River below New Bullards Bar Dam under the Proposed Action and Environmental Baseline.	8-32
8.3-2.	Difference in simulated upper tolerable water temperature exceedance probabilities for spring-run Chinook salmon adult migration and holding lifestages under the Proposed Action, relative to the Environmental Baseline.	8-33
8.3-3.	Difference in simulated upper tolerable water temperature exceedance probabilities for fall-run Chinook salmon adult migration lifestages under the Proposed Action, relative to the Environmental Baseline.	8-34

List of Tables (continued)
Description

Table No.	Description	Page No.
8.3-4.	Difference in simulated upper tolerable water temperature exceedance probabilities for spring-run Chinook salmon spawning and embryo incubation lifestages under the Proposed Action, relative to the Environmental Baseline.	8-35
8.3-5.	Difference in simulated upper tolerable water temperature exceedance probabilities for fall-run Chinook salmon spawning and embryo incubation lifestages under the Proposed Action, relative to the Environmental Baseline.	8-36
8.3-6.	Difference in simulated upper tolerable water temperature exceedance probabilities for spring-run Chinook salmon juvenile lifestages under the Proposed Action, relative to the Environmental Baseline.	8-36
8.3-7.	Difference in simulated upper tolerable water temperature exceedance probabilities for fall-run Chinook salmon juvenile lifestages under the Proposed Action, relative to the Environmental Baseline.	8-37
8.3-8.	Flows in the Middle Yuba River above the Yuba River under the Proposed Action and Environmental Baseline.	8-41
8.3-9.	Turbidity readings and locations during the sediment passage event.	8-44
8.3-10.	Flows in the Yuba River below the Middle Yuba River under the Proposed Action and Environmental Baseline.	8-49
8.3-11.	Flows in the Yuba River above New Colgate Powerhouse under the Proposed Action and Environmental Baseline.	8-50
8.3-12.	Flows in the Yuba River below New Colgate Powerhouse under the Proposed Action and Environmental Baseline.	8-51
8.3-13.	Long-term and water year type average spring-run Chinook salmon spawning WUA (percent of maximum) under the Proposed Action and Environmental Baseline.	8-57
8.3-14.	Long-term and water year type average fall-run Chinook salmon spawning WUA (percent of maximum) under the Proposed Action and Environmental Baseline.	8-59
8.3-15.	Estimated spring-run Chinook salmon redd and egg pocket potential dewatering under the Proposed Action relative to the Environmental Baseline.	8-61
8.3-16.	Estimated spring-run Chinook salmon redd and egg pocket potential dewatering under the Proposed Action relative to the Environmental Baseline for those days in the 41-year period of record during which the flow reduction criteria specified in Proposed Condition AR9 would apply.	8-63
8.3-17.	Estimated fall-run Chinook salmon redd and egg pocket potential dewatering under the Proposed Action relative to the Environmental Baseline.	8-64

List of Tables (continued)

Table No.	Description	Page No.
8.3-18.	Estimated fall-run Chinook salmon redd and egg pocket potential dewatering under the Proposed Action relative to the Environmental Baseline for those days in the 41-year period of record during which the flow reduction criteria specified in Proposed Condition AR9 would apply.....	8-65
8.3-19.	Long-term and WYT average spring-run Chinook salmon fry in-channel rearing WUA (percent of maximum) under the Proposed Action and Environmental Baseline.....	8-65
8.3-20.	Long-term and WYT average fall-run Chinook salmon fry in-channel rearing WUA (percent of maximum) under the Proposed Action and Environmental Baseline.....	8-67
8.3-21.	Long-term and WYT average spring-run Chinook salmon juvenile in-channel rearing WUA (percent of maximum) under the Proposed Action and Environmental Baseline.....	8-69
8.3-22.	Long-term and WYT average fall-run Chinook salmon juvenile in-channel rearing WUA (percent of maximum) under the Proposed Action and Environmental Baseline.....	8-71
8.3-23.	Spring-run Chinook salmon fry weighted usable area (WUA) without cover (in acres) under the Proposed Action and the Environmental Baseline over the 41-year period of evaluation for days when flows were \leq 5,000 cfs and for days when flows were $>$ 5,000 cfs, and the differences between the two scenarios over the long-term full simulation period and by water year type.....	8-73
8.3-24.	Fall-run Chinook salmon fry weighted usable area (WUA) without cover (in acres) under the Proposed Action and the Environmental Baseline over the 41-year period of evaluation for days when flows were \leq 5,000 cfs and for days when flows were $>$ 5,000 cfs, and the differences between the two scenarios over the long-term full simulation period and by water year type.....	8-75
8.3-25.	Spring-run Chinook salmon juvenile weighted usable area (WUA) without cover (in acres) under the Proposed Action and the Environmental Baseline over the 41-year period of evaluation for days when flows were \leq 5,000 cfs and for days when flows were $>$ 5,000 cfs, and the differences between the two scenarios over the long-term full simulation period and by water year type.....	8-77
8.3-26.	Fall-run Chinook salmon juvenile weighted usable area (WUA) without cover (in acres) under the Proposed Action and the Environmental Baseline over the 41-year period of evaluation for days when flows were \leq 5,000 cfs and for days when flows were $>$ 5,000 cfs, and the differences between the two scenarios over the long-term full simulation period and by water year type.....	8-79

List of Tables (continued)
Description

Table No.	Description	Page No.
8.3-27.	Difference in simulated upper tolerable water temperature exceedance probabilities for spring-run Chinook salmon lifestages under the Proposed Action relative to the Environmental Baseline.	8-84
8.3-28.	Difference in simulated upper tolerable water temperature exceedance probabilities for fall-run Chinook salmon lifestages under the Proposed Action relative to the Environmental Baseline.	8-85
8.3-29.	Allowable maximum flow reductions from the previous day average flow during the period extending from April 1 through July 15 (from Table 3 in AR9).....	8-87
8.4-1.	Exports of water from Yuba River Sub-basins from WY 1990 through 2016 averaged by water year type.	8-92
8.4-2.	Long-term average flow and average flow by water year type in the New Bullards Bar Dam Reach of the North Yuba River under the Cumulative Condition, relative to the Environmental Baseline.	8-96
8.4-3.	Long-term average flow and average flow by water year type in the Middle Yuba River above the confluence with the North Yuba River under the Cumulative Condition, relative to the Environmental Baseline.	8-97
8.4-4.	Long-term average flow and average flow by water year type in the Yuba River below the Middle Yuba River confluence under the Cumulative Condition, relative to the Environmental Baseline.	8-98
8.4-5.	Long-term average flow and average flow by water year type in the Yuba River above New Colgate Powerhouse under the Cumulative Condition, relative to the Environmental Baseline.	8-100
8.4-6.	Long-term average flow and average flow by water year type in the Yuba River below New Colgate Powerhouse under the Cumulative Condition, relative to the Environmental Baseline.	8-101
8.4-7.	Difference in simulated upper tolerable water temperature exceedance probabilities for spring-run Chinook salmon adult migration and holding lifestages under the Cumulative Condition, relative to the Environmental Baseline.....	8-102
8.4-8.	Difference in simulated upper tolerable water temperature exceedance probabilities for fall-run Chinook salmon adult migration lifestages under the Cumulative Condition, relative to the Environmental Baseline.	8-103
8.4-9.	Difference in simulated upper tolerable water temperature exceedance probabilities for spring-run Chinook salmon spawning and embryo incubation lifestages under the Cumulative Condition, relative to the Environmental Baseline.....	8-103
8.4-10.	Difference in simulated upper tolerable water temperature exceedance probabilities for fall-run Chinook salmon spawning and embryo incubation lifestages under the Cumulative Condition, relative to the Environmental Baseline.....	8-104

List of Tables (continued)

Table No.	Description	Page No.
8.4-11.	Difference in simulated upper tolerable water temperature exceedance probabilities for spring-run Chinook salmon juvenile lifestages under the Cumulative Condition, relative to the Environmental Baseline.	8-105
8.4-12.	Difference in simulated upper tolerable water temperature exceedance probabilities for fall-run Chinook salmon juvenile lifestages under the Cumulative Condition, relative to the Environmental Baseline.	8-105
8.4-13.	Long-term and water year type average spring-run Chinook salmon spawning WUA (percent of maximum) under the Cumulative Condition and Environmental Baseline.	8-107
8.4-14.	Long-term and water year type average fall-run Chinook salmon spawning WUA (percent of maximum) under the Cumulative Condition and Environmental Baseline.	8-109
8.4-15.	Estimated spring-run Chinook salmon redd and egg pocket potential dewatering under the Cumulative Condition relative to the Environmental Baseline.	8-111
8.4-16.	Estimated spring-run Chinook salmon redd and egg pocket potential dewatering under the Cumulative Condition relative to the Environmental Baseline for those days in the 33-year period of record during which the flow reduction criteria specified in Proposed Condition AR9 would apply.	8-112
8.4-17.	Estimated fall-run Chinook salmon redd and egg pocket potential dewatering under the Cumulative Condition relative to the Environmental Baseline.	8-112
8.4-18.	Estimated fall-run Chinook salmon redd and egg pocket potential dewatering under the Cumulative Condition relative to the Environmental Baseline for those days corresponding to the specific conditions during which the flow reductions specified in Proposed Condition AR9 apply.	8-113
8.4-19.	Long-term and WYT average spring-run Chinook salmon fry in-channel rearing WUA (percent of maximum) under the Cumulative Condition and Environmental Baseline.	8-114
8.4-20.	Spring-run Chinook salmon fry weighted usable area (WUA) without cover (in acres) under the Cumulative Condition and the Environmental Baseline over the 33-year period of evaluation for days when flows were \leq 5,000 cfs and for days when flows were $>$ 5,000 cfs, and the differences between the two scenarios over the long-term full simulation period and by water year type.	8-116
8.4-21.	Long-term and WYT average spring-run Chinook salmon juvenile in-channel rearing WUA (percent of maximum) under the Cumulative Condition and Environmental Baseline.	8-117

List of Tables (continued)

Table No.	Description	Page No.
8.4-22.	Spring-run Chinook salmon juvenile weighted usable area (WUA) without cover (in acres) under the Cumulative Condition and the Environmental Baseline over the 33-year period of evaluation for days when flows were $\leq 5,000$ cfs and for days when flows were $> 5,000$ cfs, and the differences between the two scenarios over the long-term full simulation period and by water year type.	8-119
8.4-23.	Long-term and WYT average fall-run Chinook salmon fry in-channel rearing WUA (percent of maximum) under the Cumulative Condition and Environmental Baseline.	8-121
8.4-24.	Fall-run Chinook salmon fry weighted usable area (WUA) without cover (in acres) under the Cumulative Condition and the Environmental Baseline over the 33-year period of evaluation for days when flows were $\leq 5,000$ cfs and for days when flows were $> 5,000$ cfs, and the differences between the two scenarios over the long-term full simulation period and by water year type.	8-122
8.4-25.	Long-term and WYT average fall-run Chinook salmon juvenile in-channel rearing WUA (percent of maximum) under the Cumulative Condition and Environmental Baseline.	8-124
8.4-26.	Fall-run Chinook salmon juvenile weighted usable area (WUA) without cover (in acres) under the Cumulative Condition and the Environmental Baseline over the 33-year period of evaluation for days when flows were $\leq 5,000$ cfs and for days when flows were $> 5,000$ cfs, and the differences between the two scenarios over the long-term full simulation period and by water year type.	8-126
8.4-27.	Difference in simulated upper tolerable water temperature exceedance probabilities for spring-run Chinook salmon lifestages under the Cumulative Condition relative to the Environmental Baseline.	8-130
8.4-28.	Difference in simulated upper tolerable water temperature exceedance probabilities for fall-run Chinook salmon lifestages under the Cumulative Condition relative to the Environmental Baseline.	8-132

List of Attachments

None.

Page Left Blank